

AD-A031 939

WISCONSIN UNIV MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

A SIMPLE PROOF OF RAMANUJAN'S SUMMATION OF THE SUB 1 PSI SUB 1.(U)

AUG 76 G E ANDREWS, R ASKEY

DAA629-75-C-0024

NL

UNCLASSIFIED

MRC-TSR-1669


1 OF 1
AD
A031939



END

DATE
FILMED

1 - 77



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

ADA031939

⑨ MRC Technical Summary Report #1669

⑥ A SIMPLE PROOF OF RAMANUJAN'S
SUMMATION OF THE ψ_1 •

⑩ George E. Andrews
Richard Askey

see tag

⑭ MRC-TSR-1669

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

⑪ Aug 1976

⑫ 8p.
(Received May 20, 1976)

⑮ DAAG29-75-C-0024,
✓ NSF-MPS-75-06687

Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P.O. Box 12211
Research Triangle Park
North Carolina 27709

and

National Science Foundation
Washington, D. C. 20550

221200

bag

UNIVERSITY OF WISCONSIN - MADISON
MATHEMATICS RESEARCH CENTER

A SIMPLE PROOF OF RAMANUJAN'S SUMMATION OF THE ${}_1\psi_1$.

George E. Andrews and Richard Askey

Technical Summary Report # 1669
August 1976


ABSTRACT

A simple proof by functional equations is given for Ramanujan's ${}_1\psi_1$ sum. Ramanujan's sum is a useful extension of Jacobi's triple product formula, and has recently become important in the treatment of certain orthogonal polynomials defined by basic hypergeometric series.

AMS(MOS) Subject Classification: 33A25, 33A30

Key Words: Basic hypergeometric functions, Theta functions,
Ramanujan sum, Jacobi triple product.

Work Unit No. 2 (Other Mathematical Methods)



Sponsored by the United States Army under Contract No. DAAG29-75-C-0024, and by the National Science Foundation under Grants 74-07282 and MPS 75-06687 A02.

A SIMPLE PROOF OF RAMANUJAN'S SUMMATION OF THE ${}_1\psi_1$

George E. Andrews⁽¹⁾ and Richard Askey⁽²⁾

In [5; p. 222, eq. (12.12.2)] G. H. Hardy alludes to Ramanujan's "... remarkable formula with many parameters.":

$$(1) \quad \sum_{n=-\infty}^{\infty} \frac{(a;q)_n x^n}{(b;q)_n} \equiv {}_1\psi_1 \left(\begin{matrix} a;q, x \\ b \end{matrix} \right) = \frac{(b/a, q)_{\infty} (q;q)_{\infty} (q/ax;q)_{\infty} (ax;q)_{\infty}}{(b;q)_{\infty} (b/ax;q)_{\infty} (q/a;q)_{\infty} (x;q)_{\infty}},$$

where $|\frac{b}{a}| < |x| < 1$, $|q| < 1$, $(a;q)_{\infty} = \prod_{n=0}^{\infty} (1-aq^n)$, and $(a;q)_n = (a;q)_{\infty} / (aq^n;q)_{\infty}$.

There are four published proofs of this result ([1], [2], [4] and [7]). Those in [1], [2] and [7] rely on somewhat tricky rearrangement of series and on the q-analog of Gauss's summation [10; p. 97, eq. (3.3.2.5)]

$$(2) \quad \sum_{n=0}^{\infty} \frac{(a;q)_n (b;q)_n (\frac{c}{ab})^n}{(c;q)_n (q;a)_n} = \frac{(c/a;q)_{\infty} (c/b;q)_{\infty}}{(c;q)_{\infty} (c/ab;q)_{\infty}},$$

where $|c| < \min(1, |ab|)$. The other proof uses the q-analogue of the binomial series [10; p. 92, eq. (3.2.2.11)]:

$$(3) \quad \sum_{n=0}^{\infty} \frac{(a;q)_n}{(q;q)_n} t^n = \frac{(at;q)_{\infty}}{(t;q)_{\infty}}, \quad |t| < 1, \quad |q| < 1,$$

but it is far from simple. Since Ramanujan's summation (1) has recently become important in the treatment of certain orthogonal polynomials defined

by basic hypergeometric series [3], it has become worthwhile to present an almost trivial proof of (1). Another very simple proof has been found by M. Ismail [6].

Proof of (1). We begin by noting that for $|q| < 1$, $f(b) \equiv {}_1\psi_1\left(\begin{smallmatrix} a;q \\ b \end{smallmatrix}; x\right)$ is an analytic function of b inside $|b| < \min(1, |ax|)$, since

$$(4) \quad f(b) = \sum_{n=0}^{\infty} \frac{(a;q)_n x^n}{(b;q)_n} + \sum_{n=1}^{\infty} \frac{(1 - \frac{b}{q}) \dots (1 - \frac{b}{q^n}) x^{-n}}{(1 - \frac{a}{q^n}) \dots (1 - \frac{a}{q})}.$$

Furthermore,

$$\begin{aligned} (5) \quad {}_1\psi_1\left(\begin{smallmatrix} a;q,x \\ b \end{smallmatrix}\right) - a {}_1\psi_1\left(\begin{smallmatrix} a;q,qx \\ b \end{smallmatrix}\right) \\ = \sum_{n=-\infty}^{\infty} \frac{(a;q)_{n+1} x^n}{(b;q)_n} = x^{-1} (1 - \frac{b}{q}) \sum_{n=-\infty}^{\infty} \frac{(a;q)_{n+1} x^{n+1}}{(\frac{b}{q};q)_{n+1}} \\ = x^{-1} (1 - \frac{b}{q}) {}_1\psi_1\left(\begin{smallmatrix} a;q,x \\ b/q \end{smallmatrix}\right). \end{aligned}$$

Hence

$$\begin{aligned} (6) \quad f(bq) - x^{-1} (1-b) f(b) &= a \sum_{n=-\infty}^{\infty} \frac{(a;q)_n q^n x^n}{(bq;q)_n} \\ &= -a b^{-1} \sum_{n=-\infty}^{\infty} \frac{(a;q)_n (1-bq^n - 1) x^n}{(bq;q)_n} = -ab^{-1} (1-b) f(b) + ab^{-1} f(bq), \end{aligned}$$

and so

$$(1 - \frac{a}{b}) f(bq) = (1-b)(x^{-1} - ab^{-1}) f(b),$$

or

$$(7) \quad f(b) = \frac{(1 - \frac{b}{a})}{(1-b)(1 - \frac{b}{ax})} f(bq).$$

If we iterate (7) $n-1$ times we find that

$$(8) \quad f(b) = \frac{(b/a; q)_n}{(b; q)_n (b/ax; q)_n} f(bq^n),$$

and since $f(b)$ is analytic in the neighborhood of 0 given by $|b| < |ax|$, we obtain in the limit as $n \rightarrow \infty$,

$$(9) \quad f(b) = \frac{(b/a; q)_\infty f(0)}{(b; q)_\infty (b/ax; q)_\infty}.$$

Now we observe from (4) and (3) that

$$(10) \quad f(q) = \sum_{n=0}^{\infty} \frac{(a; q)_n x^n}{(q; q)_n} = \frac{(ax, q)_\infty}{(x; q)_\infty}.$$

This allows us to evaluate $f(0)$ by setting $b = q$ in (9):

$$(11) \quad f(0) = \frac{(q; q)_\infty \left(\frac{q}{ax}; q\right)_\infty f(q)}{(q/a; q)_\infty} \\ = \frac{(q; q)_\infty \left(\frac{q}{ax}; q\right)_\infty (ax; q)_\infty}{(q/a; q)_\infty (x; q)_\infty}.$$

Finally we may utilize (11) to eliminate $f(0)$ from (9):

$$(12) \quad {}_1\psi_1 \left(\begin{matrix} a; q, x \\ b \end{matrix} \right) = f(b) = \frac{(b/a; q)_\infty (q; q)_\infty (q/ax; q)_\infty (ax; q)_\infty}{(b; q)_\infty (b/ax; q)_\infty (q/a, q)_\infty (x; q)_\infty},$$

as desired.

Note that Jacobi's triple product identity follows directly from (1)

if we replace a by a^{-1} , x by $z\alpha$ and then set $\alpha = b = 0$:

$$(13) \quad \sum_{n=-\infty}^{\infty} (-1)^n q^{n(n-1)/2} z^n = (q; q)_\infty (q/z; q)_\infty (z; q)_\infty.$$

I. J. Schoenberg has pointed out an interesting property of $\frac{(a;q)_n}{(b;q)_n}$ which follows from Ramanujan's sum. A sequence a_n , $n = 0, \pm 1, \dots$, is said to be totally positive if all subdeterminants of the doubly infinite matrix $A = (a_{i-j})_{-\infty < i, j < \infty}$ are nonnegative. Schoenberg [9] proved that a sequence a_n is totally positive if the bilateral generating function $f(z) = \sum_{-\infty}^{\infty} a_n z^n$ has the representation

$$(14) \quad f(z) = e^{cz+dz^{-1}} \prod_{i=1}^{\infty} \frac{(1 + \alpha_i z)(1 + \delta_i z^{-1})}{(1 - \beta_i z)(1 - \gamma_i z^{-1})},$$

$$c, d, \alpha_i, \beta_i, \gamma_i, \delta_i \geq 0, \quad \sum_{i=1}^{\infty} (\alpha_i + \beta_i + \gamma_i + \delta_i) < \infty,$$

in the interior of an annulus centered at the origin.

If $a < b < 0$ in (1) then, the generating function has the form (14) and so

$$a_n = \frac{(a;q)_n}{(b;q)_n} = \prod_{k=0}^{\infty} \frac{(1 - bq^{k+n})(1 - aq^k)}{(1 - aq^{k+n})(1 - bq^k)}$$

is a totally positive sequence for $a < b \leq 0$, $0 < q < 1$. Schoenberg [9] proved this when $b = 0$. For an extended discussion of totally positive sequences see Karlin [8].

References

1. G. E. Andrews, On Ramanujan's summation of ${}_1\psi_1(a; b; z)$, Proc. American Math. Soc., 22 (1969), 552-553.
2. G. E. Andrews, On a transformation of bilateral series with applications, Proc. American Math. Soc., 25 (1970), 554-558.
3. G. E. Andrews and R. Askey, Monograph, to appear.
4. W. Hahn, Beiträge zur Theorie der Heineschen Reihen, Math. Nach., 2 (1949), 340-379.
5. G. H. Hardy, Ramanujan, Cambridge University Press, Cambridge, 1940 (Reprinted: Chelsea, New York).
6. M. Ismail, personal communication.
7. M. Jackson, On Lerch's transcendant and the basic bilateral hypergeometric series ${}_2\psi_2$, J. London Math. Soc., 25 (1950), 189-196.
8. S. Karlin, Total Positivity, Volume One, Stanford University Press, Stanford, 1968.
9. I. J. Schoenberg, Some analytical aspects of the problem of smoothing, Studies and Essays Presented to R. Courant on his 60th Birthday, Interscience Publishers, New York, 1948, 351-370.
10. L. J. Slater, Generalized Hypergeometric Functions, Cambridge University Press, Cambridge, 1966.

Footnotes:

- (1) Partially sponsored by NSF Grant 74-07282 and by the United States Army under Contract No. DAAG29-75-C-0024.
- (2) Partially supported by NSF Grant MPS 75-06687 A02.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1669 ✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A SIMPLE PROOF OF RAMANUJAN'S SUMMATION OF THE ψ_1 ✓		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) George E. Andrews and Richard Askey		8. CONTRACT OR GRANT NUMBER(s) DAAG29-75-C-0024 ✓ MPS 75-06687 A02 74 -07282
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706 ✓		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below		12. REPORT DATE August 1976
		13. NUMBER OF PAGES 5
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U. S. Army Research Office and National Science Foundation P.O. Box 12211 Washington, D. C. 20550 Research Triangle Park, North Carolina 27709		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Basic hypergeometric functions, Theta functions, Ramanujan sum, Jacobi triple product		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A simple proof by functional equations is given for Ramanujan's ψ_1 sum. Ramanujan's sum is a useful extension of Jacobi's triple product formula, and has recently become important in the treatment of certain orthogonal polynomials defined by basic hypergeometric series. sub 1 psi sub 1		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)